

With  $K$  defined by Equation (7), the function

$$F(x) = f(x) - \phi_n(x)$$

measures the difference between the original function  $f$  and the approximating function  $\phi_n$  for each  $x$  in  $[a, b]$ .

We now use Rolle's Theorem (Section 4.2). First, because  $F(a) = F(b) = 0$  and both  $F$  and  $F'$  are continuous on  $[a, b]$ , we know that

$$F'(c_1) = 0 \quad \text{for some } c_1 \text{ in } (a, b).$$

Next, because  $F'(a) = F'(c_1) = 0$  and both  $F'$  and  $F''$  are continuous on  $[a, c_1]$ , we know that

$$F''(c_2) = 0 \quad \text{for some } c_2 \text{ in } (a, c_1).$$

Rolle's Theorem, applied successively to  $F''$ ,  $F'''$ , ...,  $F^{(n-1)}$  implies the existence of

$$\begin{aligned} c_3 & \text{ in } (a, c_2) && \text{such that } F'''(c_3) = 0, \\ c_4 & \text{ in } (a, c_3) && \text{such that } F^{(4)}(c_4) = 0, \\ & \vdots && \\ c_n & \text{ in } (a, c_{n-1}) && \text{such that } F^{(n)}(c_n) = 0. \end{aligned}$$

Finally, because  $F^{(n)}$  is continuous on  $[a, c_n]$  and differentiable on  $(a, c_n)$ , and  $F^{(n)}(a) = F^{(n)}(c_n) = 0$ , Rolle's Theorem implies that there is a number  $c_{n+1}$  in  $(a, c_n)$  such that

$$F^{(n+1)}(c_{n+1}) = 0. \tag{8}$$

If we differentiate  $F(x) = f(x) - P_n(x) - K(x-a)^{n+1}$  a total of  $n+1$  times, we get

$$F^{(n+1)}(x) = f^{(n+1)}(x) - 0 - (n+1)!K. \tag{9}$$

Equations (8) and (9) together give

$$K = \frac{f^{(n+1)}(c)}{(n+1)!} \quad \text{for some number } c = c_{n+1} \text{ in } (a, b). \tag{10}$$

Equations (7) and (10) give

$$f(b) = P_n(b) + \frac{f^{(n+1)}(c)}{(n+1)!} (b-a)^{n+1}.$$

This concludes the proof. □

**EXERCISES 11.9**

**Taylor Series by Substitution**

Use substitution (as in Example 4) to find the Taylor series at  $x = 0$  of the functions in Exercises 1-6.

- |                                       |                      |                             |
|---------------------------------------|----------------------|-----------------------------|
| 1. $e^{-5x}$                          | 2. $e^{-x/2}$        | 3. $5 \sin(-x)$             |
| 4. $\sin\left(\frac{\pi x}{2}\right)$ | 5. $\cos \sqrt{x+1}$ | 6. $\cos(x^{3/2}/\sqrt{2})$ |

**More Taylor Series**

Find Taylor series at  $x = 0$  for the functions in Exercises 7-18.

- |                                   |                    |                                 |
|-----------------------------------|--------------------|---------------------------------|
| 7. $xe^x$                         | 8. $x^2 \sin x$    | 9. $\frac{x^2}{2} - 1 + \cos x$ |
| 10. $\sin x - x + \frac{x^3}{3!}$ | 11. $x \cos \pi x$ | 12. $x^2 \cos(x^2)$             |

13.  $\cos^2 x$  (Hint:  $\cos^2 x = (1 + \cos 2x)/2$ .)  
 14.  $\sin^2 x$       15.  $\frac{x^2}{1-2x}$       16.  $x \ln(1+2x)$   
 17.  $\frac{1}{(1-x)^2}$       18.  $\frac{2}{(1-x)^3}$

### Error Estimates

19. For approximately what values of  $x$  can you replace  $\sin x$  by  $x - (x^3/6)$  with an error of magnitude no greater than  $5 \times 10^{-4}$ ? Give reasons for your answer.  
 20. If  $\cos x$  is replaced by  $1 - (x^2/2)$  and  $|x| < 0.5$ , what estimate can be made of the error? Does  $1 - (x^2/2)$  tend to be too large, or too small? Give reasons for your answer.  
 21. How close is the approximation  $\sin x = x$  when  $|x| < 10^{-3}$ ? For which of these values of  $x$  is  $x < \sin x$ ?  
 22. The estimate  $\sqrt{1+x} = 1 + (x/2)$  is used when  $x$  is small. Estimate the error when  $|x| < 0.01$ .  
 23. The approximation  $e^x = 1 + x + (x^2/2)$  is used when  $x$  is small. Use the Remainder Estimation Theorem to estimate the error when  $|x| < 0.1$ .  
 24. (Continuation of Exercise 23.) When  $x < 0$ , the series for  $e^x$  is an alternating series. Use the Alternating Series Estimation Theorem to estimate the error that results from replacing  $e^x$  by  $1 + x + (x^2/2)$  when  $-0.1 < x < 0$ . Compare your estimate with the one you obtained in Exercise 23.  
 25. Estimate the error in the approximation  $\sinh x = x + (x^3/3!)$  when  $|x| < 0.5$ . (Hint: Use  $R_4$ , not  $R_3$ .)  
 26. When  $0 \leq h \leq 0.01$ , show that  $e^h$  may be replaced by  $1 + h$  with an error of magnitude no greater than 0.6% of  $h$ . Use  $e^{0.01} = 1.01$ .  
 27. For what positive values of  $x$  can you replace  $\ln(1+x)$  by  $x$  with an error of magnitude no greater than 1% of the value of  $x$ ?  
 28. You plan to estimate  $\pi/4$  by evaluating the Maclaurin series for  $\tan^{-1} x$  at  $x = 1$ . Use the Alternating Series Estimation Theorem to determine how many terms of the series you would have to add to be sure the estimate is good to two decimal places.  
 29. a. Use the Taylor series for  $\sin x$  and the Alternating Series Estimation Theorem to show that

$$1 - \frac{x^2}{6} < \frac{\sin x}{x} < 1, \quad x \neq 0.$$

- b. Graph  $f(x) = (\sin x)/x$  together with the functions  $y = 1 - (x^2/6)$  and  $y = 1$  for  $-5 \leq x \leq 5$ . Comment on the relationships among the graphs.  
 30. a. Use the Taylor series for  $\cos x$  and the Alternating Series Estimation Theorem to show that

$$\frac{1}{2} - \frac{x^2}{24} < \frac{1 - \cos x}{x^2} < \frac{1}{2}, \quad x \neq 0.$$

(This is the inequality in Section 2.2, Exercise 52.)

- b. Graph  $f(x) = (1 - \cos x)/x^2$  together with  $y = (1/2) - (x^2/24)$  and  $y = 1/2$  for  $-9 \leq x \leq 9$ . Comment on the relationships among the graphs.

### Finding and Identifying Maclaurin Series

Recall that the Maclaurin series is just another name for the Taylor series at  $x = 0$ . Each of the series in Exercises 31–34 is the value of the Maclaurin series of a function  $f(x)$  at some point. What function and what point? What is the sum of the series?

31.  $(0.1) - \frac{(0.1)^3}{3!} + \frac{(0.1)^5}{5!} - \dots + \frac{(-1)^k(0.1)^{2k+1}}{(2k+1)!} + \dots$   
 32.  $1 - \frac{\pi^2}{4^2 \cdot 2!} + \frac{\pi^4}{4^4 \cdot 4!} - \dots + \frac{(-1)^k(\pi)^{2k}}{4^{2k} \cdot (2k!)} + \dots$   
 33.  $\frac{\pi}{3} - \frac{\pi^3}{3^3 \cdot 3} + \frac{\pi^5}{3^5 \cdot 5} - \dots + \frac{(-1)^k \pi^{2k+1}}{3^{2k+1}(2k+1)} + \dots$   
 34.  $(\pi/4) - \frac{(\pi/4)^2}{2} + \frac{(\pi/4)^3}{3} - \dots + (-1)^{k-1} \frac{(\pi/4)^k}{k} + \dots$

35. Multiply the Maclaurin series for  $e^x$  and  $\sin x$  together to find the first five nonzero terms of the Maclaurin series for  $e^x \sin x$ .  
 36. Multiply the Maclaurin series for  $e^x$  and  $\cos x$  together to find the first five nonzero terms of the Maclaurin series for  $e^x \cos x$ .  
 37. Use the identity  $\sin^2 x = (1 - \cos 2x)/2$  to obtain the Maclaurin series for  $\sin^2 x$ . Then differentiate this series to obtain the Maclaurin series for  $2 \sin x \cos x$ . Check that this is the series for  $\sin 2x$ .  
 38. (Continuation of Exercise 37.) Use the identity  $\cos^2 x = \cos 2x + \sin^2 x$  to obtain a power series for  $\cos^2 x$ .

### Theory and Examples

39. **Taylor's Theorem and the Mean Value Theorem** Explain how the Mean Value Theorem (Section 4.2, Theorem 4) is a special case of Taylor's Theorem.  
 40. **Linearizations at inflection points** Show that if the graph of a twice-differentiable function  $f(x)$  has an inflection point at  $x = a$ , then the linearization of  $f$  at  $x = a$  is also the quadratic approximation of  $f$  at  $x = a$ . This explains why tangent lines fit so well at inflection points.  
 41. **The (second) second derivative test** Use the equation

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(c_2)}{2}(x-a)^2$$

to establish the following test.

Let  $f$  have continuous first and second derivatives and suppose that  $f'(a) = 0$ . Then

- a.  $f$  has a local maximum at  $a$  if  $f'' \leq 0$  throughout an interval whose interior contains  $a$ ;  
 b.  $f$  has a local minimum at  $a$  if  $f'' \geq 0$  throughout an interval whose interior contains  $a$ .

42. **A cubic approximation** Use Taylor's formula with  $a = 0$  and  $n = 3$  to find the standard cubic approximation of  $f(x) = 1/(1-x)$  at  $x = 0$ . Give an upper bound for the magnitude of the error in the approximation when  $|x| \leq 0.1$ .

43. a. Use Taylor's formula with  $n = 2$  to find the quadratic approximation of  $f(x) = (1+x)^k$  at  $x = 0$  ( $k$  a constant).

b. If  $k = 3$ , for approximately what values of  $x$  in the interval  $[0, 1]$  will the error in the quadratic approximation be less than  $1/100$ ?

#### 44. Improving approximations to $\pi$

a. Let  $P$  be an approximation of  $\pi$  accurate to  $n$  decimals. Show that  $P + \sin P$  gives an approximation correct to  $3n$  decimals. (Hint: Let  $P = \pi + x$ .)

b. Try it with a calculator.

45. **The Taylor series generated by  $f(x) = \sum_{n=0}^{\infty} a_n x^n$  is  $\sum_{n=0}^{\infty} a_n x^n$**  A function defined by a power series  $\sum_{n=0}^{\infty} a_n x^n$  with a radius of convergence  $c > 0$  has a Taylor series that converges to the function at every point of  $(-c, c)$ . Show this by showing that the Taylor series generated by  $f(x) = \sum_{n=0}^{\infty} a_n x^n$  is the series  $\sum_{n=0}^{\infty} a_n x^n$  itself.

An immediate consequence of this is that series like

$$x \sin x = x^2 - \frac{x^4}{3!} + \frac{x^6}{5!} - \frac{x^8}{7!} + \dots$$

and

$$x^2 e^x = x^2 + x^3 + \frac{x^4}{2!} + \frac{x^5}{3!} + \dots,$$

obtained by multiplying Taylor series by powers of  $x$ , as well as series obtained by integration and differentiation of convergent power series, are themselves the Taylor series generated by the functions they represent.

46. **Taylor series for even functions and odd functions** (Continuation of Section 11.7, Exercise 45.) Suppose that  $f(x) = \sum_{n=0}^{\infty} a_n x^n$  converges for all  $x$  in an open interval  $(-c, c)$ . Show that

a. If  $f$  is even, then  $a_1 = a_3 = a_5 = \dots = 0$ , i.e., the Taylor series for  $f$  at  $x = 0$  contains only even powers of  $x$ .

b. If  $f$  is odd, then  $a_0 = a_2 = a_4 = \dots = 0$ , i.e., the Taylor series for  $f$  at  $x = 0$  contains only odd powers of  $x$ .

#### 47. Taylor polynomials of periodic functions

a. Show that every continuous periodic function  $f(x)$ ,  $-\infty < x < \infty$ , is bounded in magnitude by showing that there exists a positive constant  $M$  such that  $|f(x)| \leq M$  for all  $x$ .

b. Show that the graph of every Taylor polynomial of positive degree generated by  $f(x) = \cos x$  must eventually move away from the graph of  $\cos x$  as  $|x|$  increases. You can see this in Figure 11.13. The Taylor polynomials of  $\sin x$  behave in a similar way (Figure 11.15).

48. a. Graph the curves  $y = (1/3) - (x^2)/5$  and  $y = (x - \tan^{-1} x)/x^3$  together with the line  $y = 1/3$ .

b. Use a Taylor series to explain what you see. What is

$$\lim_{x \rightarrow 0} \frac{x - \tan^{-1} x}{x^3} ?$$

### Euler's Identity

49. Use Equation (6) to write the following powers of  $e$  in the form  $a + bi$ .

a.  $e^{-i\pi}$                       b.  $e^{i\pi/4}$                       c.  $e^{-i\pi/2}$

50. Use Equation (6) to show that

$$\cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2} \quad \text{and} \quad \sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}.$$

51. Establish the equations in Exercise 50 by combining the formal Taylor series for  $e^{i\theta}$  and  $e^{-i\theta}$ .

52. Show that

a.  $\cosh i\theta = \cos \theta$ ,                      b.  $\sinh i\theta = i \sin \theta$ .

53. By multiplying the Taylor series for  $e^x$  and  $\sin x$ , find the terms through  $x^5$  of the Taylor series for  $e^x \sin x$ . This series is the imaginary part of the series for

$$e^x \cdot e^{ix} = e^{(1+i)x}.$$

Use this fact to check your answer. For what values of  $x$  should the series for  $e^x \sin x$  converge?

54. When  $a$  and  $b$  are real, we define  $e^{(a+ib)x}$  with the equation

$$e^{(a+ib)x} = e^{ax} \cdot e^{ibx} = e^{ax} (\cos bx + i \sin bx).$$

Differentiate the right-hand side of this equation to show that

$$\frac{d}{dx} e^{(a+ib)x} = (a + ib) e^{(a+ib)x}.$$

Thus the familiar rule  $(d/dx)e^{kx} = ke^{kx}$  holds for  $k$  complex as well as real.

55. Use the definition of  $e^{i\theta}$  to show that for any real numbers  $\theta, \theta_1$ , and  $\theta_2$ ,

a.  $e^{i\theta_1} e^{i\theta_2} = e^{i(\theta_1 + \theta_2)}$ ,                      b.  $e^{-i\theta} = 1/e^{i\theta}$ .

56. Two complex numbers  $a + ib$  and  $c + id$  are equal if and only if  $a = c$  and  $b = d$ . Use this fact to evaluate

$$\int e^{ax} \cos bx \, dx \quad \text{and} \quad \int e^{ax} \sin bx \, dx$$

from

$$\int e^{(a+ib)x} \, dx = \frac{a - ib}{a^2 + b^2} e^{(a+ib)x} + C,$$

where  $C = C_1 + iC_2$  is a complex constant of integration.

25.  $\sum_{n=0}^{\infty} (-1)^n (n+1)(x-1)^n$  27.  $\sum_{n=0}^{\infty} \frac{e^2}{n!} (x-2)^n$   
 33.  $L(x) = 0, Q(x) = -x^2/2$  35.  $L(x) = 1, Q(x) = 1 + x^2/2$   
 37.  $L(x) = x, Q(x) = x$


## Section 11.9, pp. 819–822

1.  $\sum_{n=0}^{\infty} \frac{(-5x)^n}{n!} = 1 - 5x + \frac{5^2 x^2}{2!} - \frac{5^3 x^3}{3!} + \dots$   
 3.  $\sum_{n=0}^{\infty} \frac{5(-1)^n (-x)^{2n+1}}{(2n+1)!} = \sum_{n=0}^{\infty} \frac{5(-1)^{n+1} x^{2n+1}}{(2n+1)!}$   
 $= -5x + \frac{5x^3}{3!} - \frac{5x^5}{5!} + \frac{5x^7}{7!} + \dots$   
 5.  $\sum_{n=0}^{\infty} \frac{(-1)^n (x+1)^n}{(2n)!}$   
 7.  $\sum_{n=0}^{\infty} \frac{x^{n+1}}{n!} = x + x^2 + \frac{x^3}{2!} + \frac{x^4}{3!} + \frac{x^5}{4!} + \dots$   
 9.  $\sum_{n=2}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} = \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \dots$   
 11.  $x - \frac{\pi^2 x^3}{2!} + \frac{\pi^4 x^5}{4!} - \frac{\pi^6 x^7}{6!} + \dots = \sum_{n=0}^{\infty} \frac{(-1)^n \pi^{2n} x^{2n+1}}{(2n)!}$   
 13.  $1 + \sum_{n=1}^{\infty} \frac{(-1)^n (2x)^{2n}}{2 \cdot (2n)!} =$   
 $1 - \frac{(2x)^2}{2 \cdot 2!} + \frac{(2x)^4}{2 \cdot 4!} - \frac{(2x)^6}{2 \cdot 6!} + \frac{(2x)^8}{2 \cdot 8!} - \dots$   
 15.  $x^2 \sum_{n=0}^{\infty} (2x)^n = x^2 + 2x^3 + 4x^4 + \dots$   
 17.  $\sum_{n=1}^{\infty} nx^{n-1} = 1 + 2x + 3x^2 + 4x^3 + \dots$   
 19.  $|x| < (0.06)^{1/5} < 0.56968$   
 21.  $|\text{Error}| < (10^{-3})^3/6 < 1.67 \times 10^{-10}, -10^{-3} < x < 0$   
 23.  $|\text{Error}| < (30 \cdot 1)(0.1)^3/6 < 1.87 \times 10^{-4}$  25. 0.000293653  
 27.  $|x| < 0.02$  31.  $\sin x, x = 0.1; \sin(0.1)$   
 33.  $\tan^{-1} x, x = \pi/3$   
 35.  $e^x \sin x = x + x^2 + \frac{x^3}{3} - \frac{x^5}{30} - \frac{x^6}{90} \dots$   
 43. (a)  $Q(x) = 1 + kx + \frac{k(k-1)}{2} x^2$  (b) for  $0 \leq x < 100^{-1/3}$   
 49. (a)  $-1$  (b)  $(1/\sqrt{2})(1+i)$  (c)  $-i$   
 53.  $x + x^2 + \frac{1}{3} x^3 - \frac{1}{30} x^5 \dots$ ; will converge for all  $x$

9.  $1 + \frac{1}{2x} - \frac{1}{8x^2} + \frac{1}{16x^3}$   
 11.  $(1+x)^4 = 1 + 4x + 6x^2 + 4x^3 + x^4$   
 13.  $(1-2x)^3 = 1 - 6x + 12x^2 - 8x^3$   
 15.  $y = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} x^n = e^{-x}$  17.  $y = \sum_{n=1}^{\infty} (x^n/n!) = e^x - 1$   
 19.  $y = \sum_{n=2}^{\infty} (x^n/n!) = e^x - x - 1$  21.  $y = \sum_{n=0}^{\infty} \frac{x^{2n}}{2^n n!} = e^{x^2/2}$   
 23.  $y = \sum_{n=0}^{\infty} 2x^n = \frac{2}{1-x}$  25.  $y = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!} = \sinh x$   
 27.  $y = 2 + x - 2 \sum_{n=1}^{\infty} \frac{(-1)^{n+1} x^{2n}}{(2n)!}$   
 29.  $y = x - 2 \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!} - 3 \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!}$   
 31.  $y = a + bx + \frac{1}{6} x^3 - \frac{ax^4}{3 \cdot 4} - \frac{bx^5}{4 \cdot 5} - \frac{x^7}{6 \cdot 6 \cdot 7} +$   
 $\frac{ax^8}{3 \cdot 4 \cdot 7 \cdot 8} + \frac{bx^9}{4 \cdot 5 \cdot 8 \cdot 9} \dots$   
 33. 0.00267 35. 0.1 37. 0.0999444611 39. 0.100001  
 41.  $1/(13 \cdot 6!) \approx 0.00011$  43.  $\frac{x^3}{3} - \frac{x^7}{7 \cdot 3!} + \frac{x^{11}}{11 \cdot 5!}$   
 45. (a)  $\frac{x^2}{2} - \frac{x^4}{12}$   
 (b)  $\frac{x^2}{2} - \frac{x^4}{3 \cdot 4} + \frac{x^6}{5 \cdot 6} - \frac{x^8}{7 \cdot 8} + \dots + (-1)^{15} \frac{x^{32}}{31 \cdot 32}$   
 47.  $1/2$  49.  $-1/24$  51.  $1/3$  53.  $-1$  55. 2  
 59. 500 terms 61. 4 terms  
 63. (a)  $x + \frac{x^3}{6} + \frac{3x^5}{40} + \frac{5x^7}{112}$ , radius of convergence = 1  
 (b)  $\frac{\pi}{2} - x - \frac{x^3}{6} - \frac{3x^5}{40} - \frac{5x^7}{112}$   
 65.  $1 - 2x + 3x^2 - 4x^3 + \dots$  71. (c)  $3\pi/4$

## Section 11.11, pp. 838–839

1.  $f(x) = 1$



## Section 11.10, pp. 831–833

1.  $1 + \frac{x}{2} - \frac{x^2}{8} + \frac{x^3}{16}$  3.  $1 + \frac{1}{2}x + \frac{3}{8}x^2 + \frac{5}{16}x^3 + \dots$   
 5.  $1 - x + \frac{3x^2}{4} - \frac{x^3}{2}$  7.  $1 - \frac{x^3}{2} + \frac{3x^6}{8} - \frac{5x^9}{16}$