

**Proving that a function is increasing/decreasing on an interval.**

Consider the function  $f$  defined as  $f(x) = x^2 - 6x + 2$ . Completing the square yields

$$f(x) = (x - 3)^2 - 7$$

from which we see that the minimum value of the function is  $-7$  and that it occurs when  $x = 3$ . That is, the point  $(3, -7)$  is the lowest point on the function's graph.


When we say that a function  $f$  is a decreasing function of  $x$  on an interval  $(a, b)$ , we mean that as  $x$  increases while remaining between  $a$  and  $b$ ,  $f(x)$  decreases.

When we say that a function  $f$  is an increasing function of  $x$  on an interval  $(a, b)$ , we mean that as  $x$  increases while remaining between  $a$  and  $b$ ,  $f(x)$  increases.

The graph of  $f$  suggests that in the interval  $(-\infty, 3)$  as  $x$  increases,  $f(x)$  decreases; and that in the interval  $(3, \infty)$  as  $x$  increases,  $f(x)$  increases. Thus,  $f$  is decreasing on  $(-\infty, 3)$  and  $f$  is increasing on  $(3, \infty)$ .

Now, suppose we wish to prove that  $f$  defined by  $f(x) = (x - 3)^2 - 7$  is decreasing on  $(-\infty, 3)$ . Our strategy is to show that for any  $x_1, x_2$  where  $x_2 < x_1 < 3$ ,  $f(x_2) > f(x_1)$ .


Here's the geometry



The function  $f : f(x) = (x - 3)^2 - 7$  is decreasing on  $(-\infty, 3)$ .

*Proof*

Suppose  $x < 3$  and  $k > 0$ . Then  $x - k < x < 3$ . This means  $x - k$  is to the left of  $x$  and  $x$  is to the left of 3. If we can prove that  $f(x - k) > f(x)$ , then we will have shown that  $f$  is decreasing when  $x < 3$ . In other words, we will have shown that as long as we stay to the left of  $x = 3$ , the y-coordinate of a point moves down the y-axis as the x-coordinate of the point moves right on the x-axis



We must show that  $f(x - k) > f(x)$

$$f(x - k) = (x - k - 3)^2 - 7 \quad (1)$$

$$= [(x - 3) - k]^2 - 7 \quad (2)$$

$$= (x - 3)^2 - 2(x - 3)k + k^2 - 7 \quad (3)$$

$$= -2(x - 3)k + k^2 + (x - 3)^2 - 7 \quad (4)$$

$$= [-2(x - 3)k + k^2] + f(x) \quad (5)$$

$$> f(x) \quad (6)$$

$$\therefore f(x - k) > f(x)$$

□

Here is why line 6 follows from line 5.

Since

$$x < 3,$$

$$x - 3 < 3 - 3 = 0.$$

That is,

$$x - 3 < 0.$$

Thus,

$$(-2)(x - 3)k > 0$$

and

$$[-2(a - 3)k + k^2] > 0.$$

Certainly, a number greater than zero plus  $f(x)$  is greater than  $f(x)$ .