

## ■ Function

Def. A **function**  $f$  is a rule that assigns to each element  $x$  in a set  $A$  exactly one element, called  $f(x)$ , in a set  $B$ .

Def. A **function** is a set of ordered pairs such that each first element is paired with one and only one second element. That is, a relation  $F$  is a function if  $(x_1, y_1) \in F$  and  $(x_1, y_2) \in F$  implies  $y_1 = y_2$ .

Equivalently,

$$y_1 \neq y_2 \implies x_1 \neq x_2$$

$$x_1 = x_2 \implies y_1 = y_2$$

- **NB: The rule and the domain determine the function.**  $f(x) = x^2$ ,  $\mathcal{D} = \{x : x \geq 0\}$  is not the same function as  $g(x) = x^2$ ,  $\mathcal{D} = \{x : x > 0\}$ .

- **Mention vertical line test.**

## ■ Domain and Range

## ■ Algebra

Let  $f$  and  $g$  be functions with domains  $A$  and  $B$ . Then,

$$(f + g)(x) = f(x) + g(x) \quad \mathcal{D} = A \cap B$$

$$(f - g)(x) = f(x) - g(x) \quad \mathcal{D} = A \cap B$$

$$(fg)(x) = f(x)g(x) \quad \mathcal{D} = A \cap B$$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} \quad \mathcal{D} = \{A \cap B : g(x) \neq 0\}$$

## ■ Composition

Given two functions  $f$  and  $g$ , the **composite function**  $f \circ g$  is defined by  $(f \circ g)(x) = f(g(x))$ .

The domain of  $f \circ g$  is the set of all  $x$  in the domain of  $g$  such that  $g(x)$  is in the domain of  $f$ .

In general,  $f \circ g \neq g \circ f$ .

[EX1] Suppose  $f = x^2$  and  $g = x - 3$ . Then,

$$(f \circ g)(x) = f(g(x)) = f(x - 3) = (x - 3)^2$$

$$(g \circ f)(x) = g(f(x)) = g(x^2) = x^2 - 3.$$

Note that  $f \circ g \neq g \circ f$ .

## ■ 1-1 correspondence

Def. A function  $f$  with domain  $A$  and range  $B$  is called a **one-to-one function** if no two elements of  $A$  are assigned the same element in  $B$ ; that is,

$$f(x_1) \neq f(x_2) \text{ whenever } x_1 \neq x_2$$

Equivalently,

$$x_1 \neq x_2 \implies y_1 \neq y_2$$

$$y_1 = y_2 \implies x_1 = x_2$$

## ■ Mention horizontal line test.

## ■ Inverse of a function

Def. Let  $f$  be a function  $\{(x, y) : (x, y) \in f\}$  which is a one-to-one correspondence between its domain and its range. Then  $\{(y, x) : (x, y) \in f^{-1}\}$  is also a function called **the inverse of  $f$** .

NB: the inverse of  $f$  is often written  $f^{-1}$ .

## ■ Identity element for functions

Let  $f^{-1}$  be the inverse of  $f$ . Then  $f^{-1}(f(x)) = f(f^{-1}(x)) = x$ .

NB: The inverse function  $f^{-1}$  of  $f$  undoes the work of  $f$ . If  $f$  takes  $x_1 \in A$  to  $y_1 \in B$ , then  $f^{-1}$  takes  $y_1 \in B$  to  $x_1 \in A$ .  
Result:  $f^{-1}(f(x_1)) = x_1$ . You are right back where you started. The identity element is the function  $I(x) = x$ .

NB: The domain of the inverse of  $f$  is the range of  $f$ . The range of the inverse of  $f$  is the domain of  $f$ .

