

Simplify equation (2): $\frac{2x+y}{x-2y} = \frac{1}{3}$

Multiply both sides by $3(x-2y)$.

$$\begin{aligned} 3(2x+y) &= x-2y \\ 6x+3y &= x-2y \\ 5x+5y &= 0 \\ x+y &= 0 \end{aligned} \quad (4)$$

Equations (3) and (4), which are equivalent forms of equations (1) and (2), may now be solved by either of the standard methods.

Exercises [A-1]

Solve the following systems of equations:

1. $\begin{cases} x+4y=1 \\ x+7y=4 \end{cases}$

5. $\begin{cases} 5x+4y=3 \\ 2x+3y=4 \end{cases}$

9. $\begin{cases} .1x+3y=2.6 \\ x-1.6y=10.2 \end{cases}$

2. $\begin{cases} y=4x-7 \\ 16x-5y=25 \end{cases}$

6. $\begin{cases} x+y=14 \\ \frac{x}{y}=\frac{3}{4} \end{cases}$

10. $\begin{cases} 3x-2y+4=0 \\ \frac{x-y}{x+y}=3 \end{cases}$

3. $\begin{cases} 5x+3y=6 \\ 7x-y=11 \end{cases}$

7. $\begin{cases} \frac{1}{2}x-\frac{1}{3}y=2 \\ \frac{1}{3}x+\frac{1}{2}y=\frac{17}{3} \end{cases}$

11. $\begin{cases} y=3+.8x \\ 4y-2.2x=2 \end{cases}$

4. $\begin{cases} 5x-3y=7 \\ 15x-9y=3 \end{cases}$

8. $\begin{cases} .5x-y=1.7 \\ .3x+.4y=1.3 \end{cases}$

12. $\begin{cases} \frac{1}{u}+\frac{1}{t}=5 \\ \frac{1}{u}-\frac{1}{t}=1 \end{cases}$

14. $\begin{cases} \frac{h}{x}=.35 \\ \frac{h}{x+10}=.25 \end{cases}$

16. $\begin{cases} \frac{1}{2}(x-2)-\frac{1}{3}(y+3)=-2 \\ \frac{3(x-2)}{4}-\frac{5(y+1)}{2}=2 \end{cases}$

13. $\begin{cases} \frac{6}{x}+\frac{4}{y}=9 \\ \frac{12}{x}-\frac{2}{y}=3 \end{cases}$

15. $\begin{cases} \frac{10}{x}+3y=2 \\ \frac{8}{x}-2y=6 \end{cases}$

17. $\begin{cases} \frac{4}{3x}+\frac{1}{2y}=1 \\ \frac{5}{3x}-\frac{3}{4y}=4 \end{cases}$

18. How many pairs of values of x and y satisfy the equation $x+y=8$,
 (a) If there is no restriction on the values of the variables?
 (b) If both x and y must be positive, odd integers?
 (c) If the values of x and y must be equal?
19. If $y=7u+2$, and $u=4-3x$, eliminate u by substitution and express y in terms of x . Find the value of y for which $x=2$
20. In the equation $\frac{1}{2}x+\frac{1}{3}(y+8)=4$, find the value of y if (a) the value of x is $-\frac{5}{2}$, (b) the values of x and y are in the ratio $2:1$.
21. If the equation $3x+ky=2$ is satisfied by the pair of values $x=2$, $y=-1$, find the value of k .

Exercises [A-2]

Solve the following systems of equations:

1. $\begin{cases} 3x - y = 5 \\ 5x - y = 7 \end{cases}$
2. $\begin{cases} y = 11 - 3x \\ 5x - 3y = 2 \end{cases}$
3. $\begin{cases} 3x + 5y = 7 \\ x - 3y = 0 \end{cases}$
4. $\begin{cases} 2x + y = 4 \\ 4x + 2y = 6 \end{cases}$
5. $\begin{cases} 3x + 4y = 1 \\ 5x + 3y = 9 \end{cases}$
6. $\begin{cases} x - y = 1 \\ y = \frac{1}{2} \end{cases}$
7. $\begin{cases} \frac{2x}{3} + \frac{y}{9} = 6 \\ \frac{1}{4}x + \frac{1}{2}y = 5 \end{cases}$
8. $\begin{cases} .5x + 1.2y = 1.4 \\ .6x - 7.0y = 5.9 \end{cases}$
9. $\begin{cases} .5x + .2y = 1.65 \\ .7x - .3y = 2.6 \end{cases}$
10. $\begin{cases} 2x - 3y - 3 = 0 \\ \frac{x+y}{x-y} = \frac{1}{2} \end{cases}$
11. $\begin{cases} x = 2 - .25y \\ 9x + 1.25y = 15 \end{cases}$
12. $\begin{cases} \frac{2}{u} + \frac{1}{t} = 7 \\ \frac{2}{u} - \frac{1}{t} = 1 \end{cases}$
13. $\begin{cases} \frac{3}{x} + \frac{4}{y} = 8 \\ \frac{6}{x} - \frac{3}{y} = 5 \end{cases}$
14. $\begin{cases} \frac{h}{x} = .45 \\ \frac{h}{x+50} = .30 \end{cases}$
15. $\begin{cases} \frac{8}{x} + 3y = 5 \\ \frac{6}{x} + 2y = 3 \end{cases}$
16. $\begin{cases} \frac{1}{3}(x+1) - \frac{1}{2}(3y-1) = 1 \\ \frac{3(2x-1)}{5} - \frac{2(y-3)}{3} = 2 \end{cases}$
17. $\begin{cases} \frac{3}{2x} - \frac{1}{3y} = 2 \\ \frac{3}{4x} + \frac{5}{6y} = -2 \end{cases}$

18. How many pairs of values of x and y satisfy the equation $x + y = 10$,
 - (a) If there is no restriction on the values of the variables?
 - (b) If both x and y must be even integers, greater than zero?
 - (c) If the value of x must be 25% of the value of y ?
19. If $x = 4t + 3$, and $t = 3 - 2u$, eliminate t by substitution and express x in terms of u . Find the value of u for which $x = 0$.
20. In the equation $\frac{1}{3}(x-2) + \frac{1}{4}y = \frac{1}{2}$, find the value of x if (a) the value of y is $-\frac{2}{3}$, (b) the values of x and y are equal.
21. If the equation $4x - 3y = c$ is satisfied by the pair of values $x = \frac{1}{2}$, $y = -1$, find the value of c .
22. Show that the equations $5x - y = 21$, $x + 3y = 1$, $13x + 7y = 45$ have a common solution.

Exercises [B]

1. Show that the equation $\frac{1}{2}x + \frac{1}{3}(y-3) = \frac{1}{3}x + \frac{1}{2}(y-2)$ is equivalent to the equation $x = y$.
2. State the number of solutions that may be found for each of the following pairs of equations:

(a) $2x + 5y = 8$	(b) $2x + 5y = 8$	(c) $2x + 5y = 8$
$4x + 10y = 12$	$4x + 10y = 16$	$2x - y = 2$

3. In general, how many solutions may be found to a problem in two variables,
- If only one equation in the two variables is to be satisfied?
 - If two independent equations in the variables are to be satisfied?
 - If three independent equations in the variables are to be satisfied?
4. If $7x + 3y = 18$, find the smallest integral value of y for which the corresponding value of x is less than 1.
5. Solve for x and y : $x + y = 28$, $\frac{7}{x+y} = \frac{4}{x-y}$.
6. Show that no pair of values of x and y satisfies the three equations $3x - 2y = 8$, $5x - 4y = -3$, $\frac{x}{y} = \frac{3}{4}$.
7. Find the value that a must have if the three equations $3x - 2y = 8$, $2x + 3y = 1$, $ax + 5y = -1$ are to have a common solution.
8. Solve for x and y : $x - \frac{1}{3}y = 3y - \frac{1}{2}x = 8\frac{1}{2}$.
9. State the number of solutions that may be found for each of the following pairs of equations:
- | | | |
|-------------------|-------------------------|------------------------|
| (a) $4x - 3y = 6$ | (b) $4x - 3y = 6$ | (c) $4x - 3y = 8$ |
| $8x - 6y = 9$ | $2x - \frac{3}{2}y = 3$ | $x + \frac{3}{2}y = 4$ |
10. How many pairs of numbers may be found to satisfy the following conditions?
- The sum of the two numbers is 25.
 - The sum of the two numbers is 25, and the difference of the numbers is 12.
 - The sum of the two numbers is 25, their difference is 12, and the numbers are equal.
11. If $3x + 2y = 7$, find the smallest integral value of x for which the corresponding value of y is negative.
12. Solve for t and u :
$$\begin{cases} t - u = 5 \\ \frac{4}{u+t} = \frac{10}{u-t} \end{cases}$$
13. Show that no values of x and y satisfy the three equations $x - y = 1$, $2x - 3y = 2$, $y = \frac{1}{2}x$.
14. Find the value that b must have if the three equations $2x - y = 5$, $4x + 3y = -5$, $3x + by = 9$ are to have a common solution.
15. Solve for x and y : $\frac{1}{2}x - 7y - 5 = \frac{3}{2}x - y = 0$.

4 • SECOND COURSE IN ALGEBRA

3. $\{x \mid x = \pm 2\}$, or $\{x \mid x^2 = 4\}$ (other answers possible).
4. a. $\{B, D\}$. b. \emptyset .
5. a. $\{1, 2, 3, 4\}$. b. $\{2, -2\}$. c. None.
6. $\{0, 1, 2, 3, 4, 5, 6\}$.
7. $\{x \mid x \text{ is an integer and } 0 < x < 100\}$.
8. R is the set of rational numbers (or whatever universal set is in use).

Page 31

- | | |
|--|---|
| <u>1.</u> $(-3, 1)$ | <u>13.</u> $(2, \frac{2}{3})$ |
| <u>2.</u> $(2\frac{1}{2}, 3)$ | <u>14.</u> $(x, h) = (25, 8.75)$ |
| <u>3.</u> $(1\frac{1}{2}, -\frac{1}{2})$ | <u>15.</u> $(2, -1)$ |
| <u>4.</u> No solution | <u>16.</u> $(-2, -3)$ |
| <u>5.</u> $(-1, 2)$ | <u>17.</u> $(\frac{2}{3}, -\frac{1}{2})$ |
| <u>6.</u> $(6, 8)$ | <u>18.</u> a. An unlimited number |
| <u>7.</u> $(8, 6)$ | b. Four: $(1, 7), (3, 5), (5, 3), (7, 1)$ |
| <u>8.</u> $(3.96, 0.28)$ | c. One: $(4, 4)$ |
| <u>9.</u> $(11, 0.5)$ | <u>19.</u> $y = 30 - 21x; y = -12$ |
| <u>10.</u> $(-1, \frac{1}{2})$ | <u>20.</u> a. $y = 7\frac{3}{4}$ |
| <u>11.</u> $(-10, -5)$ | b. $y = 1$ |
| <u>12.</u> $(u, t) = (\frac{1}{3}, \frac{1}{2})$ | <u>21.</u> $k = 4$ |

Page 32

- | | |
|--|--|
| <u>1.</u> $(1, -2)$ | <u>11.</u> $(1.25, 3)$ |
| <u>2.</u> $(2\frac{1}{2}, 3\frac{1}{2})$ | <u>12.</u> $(u, t) = (\frac{1}{2}, \frac{1}{3})$ |
| <u>3.</u> $(\frac{3}{2}, \frac{1}{2})$ | <u>13.</u> $(\frac{3}{4}, 1)$ |
| <u>4.</u> No solution | <u>14.</u> $(x, h) = (100, 45)$ |
| <u>5.</u> $(3, -2)$ | <u>15.</u> $(-2, 3)$ |
| <u>6.</u> $(2, 1)$ | <u>16.</u> $(\frac{1}{2}, 0)$ |
| <u>7.</u> $(8, 6)$ | <u>17.</u> $(\frac{3}{2}, -\frac{1}{3})$ |
| <u>8.</u> $(4, -\frac{1}{2})$ | <u>18.</u> a. An unlimited number |
| <u>9.</u> $(3.5, -0.5)$ | b. Four: $(2, 8), (4, 6), (6, 4), (8, 2)$ |
| <u>10.</u> $(1, -\frac{1}{3})$ | c. One: $(2, 8)$ |

19. $x = 15 - 8u$; $u = 1\frac{7}{8}$ 21. $c = 5$
 20. a. $x = 4$ b. $x = 2$ 22. $(4, -1)$ is common sol.

Pages 32-33

2. a. None 9. a. None
 b. Unlimited number b. Unlimited number
 c. One c. One
 3. a. Unlimited number 10. a. Unlimited number
 b. One b. One: $(18\frac{1}{2}, 6\frac{1}{2})$
 c. None c. None
 4. $y = 4$ 11. $x = 3$
 5. $(22, 6)$ 12. $(t, u) = (1\frac{1}{2}, -3\frac{1}{2})$
 7. $a = 2$ 14. $b = -2$
 8. $(10, 4\frac{1}{2})$ 15. $(-\frac{1}{2}, -\frac{3}{4})$

Pages 35-36

1. $(1, 5, 3)$ 3. $(2, -1, 3)$ 5. $(r, s, t) = (9, 11, 8)$
 2. $(1, 1, 0)$ 4. $(\frac{1}{2}, -\frac{1}{4}, \frac{1}{3})$ 6. $(-\frac{1}{3}, \frac{1}{5}, \frac{1}{2})$

Page 36 (middle)

1. $(3, 2, 1)$ 3. $(-1, 4, 8)$ 5. $(s, t, u) = (6, 8, 5)$
 2. $(r, s, t) = (2, 0, 1)$ 4. $(\frac{2}{3}, -\frac{1}{3}, \frac{1}{2})$ 6. $(\frac{1}{4}, -\frac{1}{3}, \frac{1}{2})$

Page 36 (bottom)

2. $(a, b, c) = (8, 4, 3)$ 4. $(a, b, c) = (3k, 2k, 4k)$, k any constant.

Pages 40-42

1. a. $d = 47\frac{1}{2} - 6x$ 4. a. Width, $(x + 4)$ in.
 b. 43 Length, $(x + 7)$ in.
 c. $1\frac{1}{2}$ b. $(8x + 28)$ sq.in.
 2. a. hours c. 18
 b. $n = 5 + \frac{5}{24}x$ 5. a. 84 cents
 c. $6\frac{1}{4}$ b. $(90 - \frac{3}{5}x)$ cents
 d. 8 c. 35
 3. a. 35 m.p.h. 6. a. $I = 275 - 0.03x$
 b. $v = \frac{7x}{4}$ b. \$230
 c. 22.9 c. \$2700